




Gonorrhoea

Overview

- Gonorrhoea is most commonly diagnosed in [men who have sex with men, young \(heterosexual\), Aboriginal and Torres Strait Islander people](#) living in remote and very remote areas, and travellers returning from high prevalence areas overseas.
- Increasing prevalence in general population, especially in women of reproductive age.
- Previous infection does not provide immunity to new infection.
- Reduced susceptibility to first-line treatment is emerging in urban Australia and is being monitored closely.



Cause

- Neisseria gonorrhoeae, a Gram-negative intracellular diplococcus bacterium.

Clinical presentation

Anal and pharyngeal gonorrhoea infections are asymptomatic in most people, vaginal gonorrhoea is often asymptomatic (in 80% of cases), and penile urethral gonorrhoea is almost always symptomatic (in 85–90% of cases).

Symptoms

- [Penile urethral discharge](#)
- [Dysuria](#)
- [Vaginal discharge](#)
- Dyspareunia with cervicitis
- Conjunctivitis: purulent, sight-threatening
- [Anorectal symptoms](#): discharge, irritation, painful defecation, disturbed bowel function

Complications

- [Epididymo-orchitis](#) (uncommon): painful, red swollen testicle/s
- Prostatitis (very rarely)
- [Pelvic inflammatory disease \(PID\)](#): dyspareunia, intermenstrual bleeding, post-coital bleeding, discharge
- Bartholin gland abscess
- Disseminated disease (rarely):
 - macular rash that may include necrotic pustules
 - septic arthritis
- Meningitis or endocarditis (rarely)

See [STI Atlas](#) for images.

Diagnosis

ALWAYS collect samples for gonococcal culture before treating gonorrhoea, to determine antimicrobial sensitivity and contribute to antimicrobial resistance surveillance. However, do not delay treatment to wait for these culture results.

Site/Specimen	Test	Consideration
First pass urine (FPU)	NAAT	<p>In people who do not have a vagina or if endocervical swab/self-collected vaginal swab cannot be taken</p> <p>Less sensitive than self-collected vaginal swab</p> <p>For men who have sex with men, also collect anal and pharyngeal swabs for NAAT even if asymptomatic at these sites</p>

		pharyngeal swab for NAAT even if asymptomatic at these sites
Penile urethral swab	Culture	Only required if discharge or other local symptoms present, or before administering antibiotics If men who have sex with men , also collect anal and pharyngeal swab for NAAT even if asymptomatic at these sites
Clinician-collected endocervical swab	NAAT +/- culture	Best test if examined In asymptomatic patients, an endocervical swab for NAAT is sufficient. However, if the patient has vaginal symptoms, collect additional swab for gonococcal culture
Self-collected vaginal swab	NAAT +/- culture	Best test if not examined In asymptomatic patients, a vaginal swab for NAAT is sufficient. However, if the patient has vaginal symptoms, collect additional swab for gonococcal culture
Anorectal swab	NAAT +/- culture	Collect for all men who have sex with men , and any patient with anorectal symptoms In asymptomatic patients, a self-collected or practitioner-collected rectal swab for NAAT is sufficient. However, if the patient has anorectal symptoms, collect additional swab for gonococcal culture
Pharyngeal swab	NAAT +/- culture	Collect for all men who have sex with men , and for anyone else with multiple sexual partners.

NAAT – Nucleic acid amplification test.

Gonococcal culture should always be collected before antibiotics are administered, but treatment should be administered without waiting for culture results.

Specimen collection guidance

[Clinician collected](#) | [Self-collection](#)

Investigations

- NAATs are highly sensitive, allow for patient self-sampling and can be used in non-clinical and non-urban settings. They are not validated for non-genital sites however, and rarely false positives can occur. NAATs are the most common gonorrhoea test used in Australia.
- Gonococcal culture has high specificity and allows for antibiotic susceptibility testing but is much less sensitive than NAAT. Culture samples should be obtained from all infected sites at the time of treatment to determine antibiotic susceptibility. Culture accuracy depends on stringent incubation and transport conditions and should reach the laboratory within 24 hours. Clinicians must specifically request 'gonococcal culture' rather than general 'culture', as gonococci require specific culture conditions.

Management

Gonococcal culture samples should always be collected from all infected sites before antibiotics are administered, but treatment should be administered without waiting for culture results.

Principal treatment options		
Situation	Recommended	Alternative
Uncomplicated genital and anorectal infection	Ceftriaxone 500 mg IMI, stat. in 2 mL 1% lignocaine PLUS Azithromycin 1 g PO, stat.	Alternative treatments are not recommended because of high levels of resistance, EXCEPT for some remote Australian locations and severe allergic reactions. Seek local specialist advice.
Uncomplicated pharyngeal infection*	Ceftriaxone 500 mg IMI, stat. in 2 mL 1% lignocaine PLUS Azithromycin 2 g PO, stat.	Alternative treatments are not recommended because of high levels of resistance, EXCEPT for some remote Australian locations and severe allergic reactions. Azithromycin given as 1 g followed by 1 g 6 hours later may reduce gastrointestinal side-effects.
Adult gonococcal conjunctivitis	Ceftriaxone 1 g IMI, stat. in 2 mL 1% lignocaine PLUS Azithromycin 1 g PO, stat.	Alternative treatments are not recommended because of high levels of resistance, EXCEPT for some remote Australian locations and severe allergic reactions.

BD: twice a day

IMI: intramuscular injection

PO: orally

stat.: immediately

*If a patient has received the recommended treatment for genital or anorectal gonorrhoea at the time of testing, and if they're found to also have pharyngeal gonorrhoea, they do not need to be re-treated with the higher dose of azithromycin, but a test of cure is recommended.

Treatment advice

- Reduced susceptibility to the first-line treatment of intramuscular injection ceftriaxone and azithromycin is emerging in urban Australia.
- Sharing of antimicrobial resistance genetic material between bacteria and reduced drug penetration to pharyngeal mucosa makes it the most likely site of treatment failure.
- Dual antibiotic treatment is recommended to create a pharmacological barrier to the development of more widespread resistance to treatment.
- Ceftriaxone monotherapy should not be used outside specialist centres monitoring culture-based antibiotic susceptibility.
- Ciprofloxacin 500 mg orally immediately can be used to treat gonorrhoea, but only if susceptibility has been confirmed on NAAT or culture, and this should not delay treatment.
- If a patient has an intrauterine device (IUD), leave it in place and treat as recommended. Seek specialist advice as needed.

Other immediate management

- Advise no sexual contact for **7 days** after treatment is administered
- Advise no sex with partners from the last **2 months** until the partners have been tested and treated if necessary.
- Recommend partner notification.
- Provide patient with [factsheet](#).
- Notify the state or territory health department.
- Consider testing for [other sexually transmitted infections \(STIs\)](#), if not undertaken at first presentation.
- Consider human immunodeficiency virus ([HIV](#)) pre-exposure prophylaxis (PrEP) for anyone diagnosed with gonorrhoea.

Special Treatment Situations

Special considerations

- Consider seeking specialist advice before treating any complicated presentation.

Situation	Recommended
Rectal co-infection	For rectal co-infection with chlamydia , treatment should be given for gonorrhoea AND chlamydia i.e. Ceftriaxone 500 mg IMI, stat. in 2 mL 1% lignocaine PLUS Doxycycline 100 mg PO, BD 7 days if asymptomatic, but 21 days if symptomatic (see anorectal syndromes)
Pregnancy	Same as principal treatment option.
Allergy to principal treatment choice	Seek specialist advice.
Regional/remote areas where gonorrhoea is known to be penicillin susceptible. <i>To confirm check with local public health unit</i>	Amoxicillin 3 g PO, stat. PLUS Probenecid 1 g PO, stat. PLUS Azithromycin 1 g PO, stat. (when chlamydia not excluded). If the infection is likely to have been acquired beyond local or other remote locations, use principal treatment option.

Contact Tracing

- Contact tracing for gonorrhoea is a high priority and should be performed in all patients with confirmed infection.

- Contact tracing is important to prevent re-infection and reduce transmission.
- All partners should be traced back for a minimum of **2 months**.
- The diagnosing doctor is responsible for initiating and documenting a discussion about contact tracing.
- Offer testing of exposed anatomical sites to all sexual contacts.
- Consider presumptive treatment if there has been sexual contact within the past 2 weeks or when the person's individual circumstances mean later treatment may not occur.

See [Australasian Contract Tracing Manual – Gonorrhoea](#) for more information.

Follow Up

Review in **1 week** provides an opportunity to:

- Confirm patient adherence with treatment and assess for symptom resolution
- Confirm contact tracing has been undertaken or offer more contact tracing support
- Provide further sexual health education and prevention counselling, including discussion and provision of HIV PrEP.

Test of cure

For each site of infection (pharyngeal, anal or cervical), TOC by NAAT should be performed **2 weeks** after treatment is completed, especially if no gonococcal culture swab was collected before treatment, or if the culture swab indicated antimicrobial resistance, or if the patient was treated with a non-standard regimen.

Test for re-infection:

- Re-infection is common.
- Retesting at **3 months** is recommended to detect re-infection.
- Consider testing for [other sexually transmitted infections \(STIs\)](#) including Syphilis and HIV if not undertaken at first presentation or retesting after the window period.

If test of cure or retesting is positive, seek specialist advice.

Auditable Outcomes

- 100% of patients diagnosed with gonorrhoea are treated with an appropriate antibiotic regimen.
- 100% of patients are advised to avoid sexual contact for 7 days after treatment is administered.
- 100% of patients diagnosed with gonorrhoea are recalled for repeat testing after 3 months.

Endorsement: These guidelines have been endorsed by the Blood Borne Viruses and Sexually Transmitted Infections Standing Committee (BBVSS).

Developed by: the [Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine \(ASHM\)](#), ABN 48 264 545 457 | CFN 17788

Funded by: The Australian Government Department of Health

Disclaimer: Whilst the Australian Department of Health provides financial assistance to ASHM, the material contained in this resource produced by ASHM should not be taken to represent the views of the Australian Department of Health. The content of this resource is the sole responsibility of ASHM. www.ashm.org.au

Last updated: December, 2021

© ASHM, Australia, 2022